

Recovery of the immune system after exercise

Peake, Jonathan M; Neubauer, Oliver; Walsh, Neil P; Simpson, Richard J

Journal of Applied Physiology

DOI:

[10.1152/jappphysiol.00622.2016](https://doi.org/10.1152/jappphysiol.00622.2016)

Published: 01/05/2017

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Peake, J. M., Neubauer, O., Walsh, N. P., & Simpson, R. J. (2017). Recovery of the immune system after exercise. *Journal of Applied Physiology*, 122(5), 1077-1087.
<https://doi.org/10.1152/jappphysiol.00622.2016>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Recovery of the immune system after exercise

Jonathan M. Peake^{1,2}, Oliver Neubauer¹, Neil P. Walsh³, Richard J. Simpson⁴

¹ School of Biomedical Sciences and Institute of Health and Biomedical Innovation,
Queensland University of Technology, Brisbane, Australia.

² Centre of Excellence for Applied Sport Science Research, Queensland Academy of
Sport, Brisbane, Australia.

³ Extremes Research Group, School of Sport, Health and Exercise Sciences, Bangor
University, Bangor, UK.

⁴ Laboratory of Integrated Physiology, Department of Health and Human Performance,
University of Houston, Houston, Texas, USA.

Running head: Immune system recovery after exercise

Word count in main document: 5,253

Corresponding author:

Jonathan Peake
Institute of Health and Biomedical Innovation
Queensland University of Technology
Kelvin Grove, QLD 4059
Australia
Email: jonathan.peake@qut.edu.au
Phone: +61 7 3138 6140

1 ABSTRACT

2 The notion that prolonged, intense exercise causes an 'open window' of
3 immunodepression during recovery after exercise is well accepted. Repeated exercise bouts
4 or intensified training without sufficient recovery may increase the risk of illness. However,
5 except for salivary IgA, clear and consistent markers of this immunodepression remain
6 elusive. Exercise increases circulating neutrophil and monocyte counts, and reduces
7 circulating lymphocyte count during recovery. This lymphopenia results from preferential
8 egress of lymphocyte subtypes with potent effector functions (e.g., NK cells, $\gamma\delta$ T cells and
9 CD8⁺ T cells). These lymphocytes most likely translocate to peripheral sites of potential
10 antigen encounter (e.g., lungs, gut). This redeployment of effector lymphocytes is an integral
11 part of the physiological stress response to exercise. Current knowledge about changes in
12 immune function during recovery from exercise is derived from assessment at the cell
13 population level of isolated cells ex vivo or in blood. This assessment can be biased by large
14 changes in the distribution of immune cells between blood and peripheral tissues during and
15 after exercise. Some evidence suggests that reduced immune cell function in vitro may
16 coincide with changes in vivo and rates of illness after exercise, but more work is required to
17 substantiate this notion. Among the various nutritional strategies and physical therapies that
18 athletes use to recover from exercise, carbohydrate supplementation is the most effective for
19 minimizing immune disturbances during exercise recovery. Sleep is an important aspect of
20 recovery, but more research is needed to determine how sleep disruption influences the
21 immune system of athletes.

22

23 **Keywords:** open window; repeated exercise bouts; immunodepression; overreaching; sleep

24 INTRODUCTION

25 The immune system is integral to the body's defense against infection. It also influences
26 other physiological systems and processes, including tissue repair, metabolism,
27 thermoregulation, sleep/fatigue, and mental health. Over the past 40 years, exercise
28 immunology has developed into its own discipline based on the recognition that the immune
29 system mediates many exercise effects and that stress responses mediated through the
30 nervous and endocrine systems play a key role in determining exercise-induced immune
31 changes (84). A classic paradigm in exercise immunology is that an 'open window' of
32 immunodepression can occur during recovery from intense exercise. In particular, this
33 paradigm proposes that after intense exercise, some immune variables (e.g., lymphocyte and
34 natural killer cell numbers, antibody production) transiently decrease below preexercise
35 levels. As a result of this immunodepression microbial agents, especially viruses, may invade
36 the host or reactivate from a latent state, leading to infection and illness (87). If exercise is
37 repeated again while the immune system is still depressed, this could lead to a greater degree
38 of immunodepression and potentially a longer window of opportunity for infection (87).

39 Exercise-induced fatigue exists on a continuum. Repeated bouts of intense exercise on the
40 same day or over several days may cause acute fatigue, as indicated by an inability to maintain
41 exercise workloads (64). An athlete who trains intensely for 1–2 weeks may experience a state
42 of 'functional overreaching', which is associated with a temporary performance decrement,
43 followed by improved performance. Intense training over an extended period without
44 sufficient balance between training and recovery may lead to 'nonfunctional overreaching'
45 (NFOR) (64). This condition is typically characterized by persistent fatigue, performance
46 decrement, muscle soreness, and psychological and hormonal disturbances that can last for

47 weeks or months. Depending on the time needed to recover from NFOR, an athlete may be
48 diagnosed (retrospectively) as experiencing ‘overtraining syndrome’ (64).

49 Recognition of the link between excessive training and risk of illness has stimulated
50 interest in nutritional supplements and physical therapies to counteract immunodepression
51 and to restore immune function after exercise training. In this mini-review, we update the
52 current state of knowledge about the temporal changes in the immune system following
53 exercise; how repeated bouts of exercise on the same day, extended periods of intense
54 training, and sleep disruption influence the immune system; and the efficacy of various
55 strategies for restoring immune function after exercise.

56

57 **LEUKOCYTE REDEPLOYMENT DURING EXERCISE AND RECOVERY**

58 A single exercise bout causes profound changes in the number and composition of blood
59 leukocytes that may persist long into exercise recovery. All major leukocyte subpopulations
60 tend to increase in number during exercise as a result of hemodynamic shear stress and/or
61 catecholamines acting on leukocyte β_2 adrenergic receptors (126). The postexercise recovery
62 period is marked by opposite effects on blood neutrophil and lymphocyte numbers.
63 Neutrophil number (and, consequently, the total leukocyte count) often continues to increase
64 long into the recovery period (up to 6 h after exercise cessation), particularly if the exercise
65 bout is prolonged (>2 h) (86). This sustained ‘neutrophilia’ is characterized by an increased
66 presence of immature, less differentiated, precursor neutrophils in the blood (117), most
67 likely in response to the increased plasma levels of soluble agents including glucocorticoids,
68 growth hormone, and cytokines such as IL-6 and granulocyte colony-stimulating factor, which

69 mobilize myeloid cells from the bone marrow (117). Although this neutrophilia following
 70 prolonged exercise is akin to that observed during bacterial infection ($>7.0 \times 10^6/\text{ml}$), 24 h of
 71 recovery is usually sufficient for neutrophil number to return to normal (126). A delayed
 72 monocytosis is sometimes observed within 1–2 h after very prolonged exercise, but monocyte
 73 number typically returns to the resting level within 6 h after exercise cessation (126).

74 By contrast, lymphocyte number decreases rapidly after exercise. Following prolonged
 75 and/or high-intensity exercise in particular, lymphocyte number commonly decreases to
 76 below the preexercise value within as little as 30 min (126). This ‘lymphopenia’ can often
 77 reach levels typical of clinical lymphopenia ($<1.0 \times 10^6/\text{ml}$) but the lymphocyte count is usually
 78 restored to both the resting and clinically normal level within 4–6 h of recovery (126). After
 79 prolonged bouts of exercise (e.g., 2 h cycling), natural killer (NK) cells (which account for most
 80 of the exercise-induced lymphocytosis) may be ~40% lower than the baseline value for up to
 81 7 d after exercise (104). Exercise-induced lymphopenia reflects the preferential movement of
 82 lymphocyte subtypes with potent effector functions (e.g., NK cells, $\gamma\delta$ T cells, and CD8^+ T cells)
 83 out of the blood. Even within these subsets, there is a preferential egress of discrete subtypes
 84 of highly differentiated NK-cells, $\gamma\delta$ T cells, and CD8^+ T cells with phenotypes associated with
 85 tissue-migrating potential, and effector capabilities (107).

86 The rapid lymphopenia observed during the early stage of exercise recovery was initially
 87 of concern, particularly because early studies reported large rates of lymphocyte apoptosis
 88 (programmed cell death) after exhaustive exercise (62). However, these findings have not
 89 been substantiated. Subsequent studies have reported lymphocyte apoptosis in the order of
 90 0–2% after exercise, even though the blood lymphocyte count was up to 30–40% lower than
 91 at rest (66, 105). Lymphocytes and monocytes leave the blood in large numbers during

92 exercise recovery under the influence of glucocorticoids. Lymphocyte subtypes that
93 preferentially egress the peripheral blood during exercise recovery also have phenotypes
94 consistent with tissue migration (e.g., expression of surface adhesion molecules, chemokine
95 receptors) (108). These lymphocytes most likely translocate to peripheral sites of potential
96 antigen encounter, such as the lungs or the gut (48).

97 The skin has long been considered a likely destination for effector lymphocytes in
98 response to exercise and stress in general (24). However, recent evidence indicates that CD8⁺
99 T cells and NK cells mobilized by exercise do not express cutaneous homing receptors on their
100 surface (121). Exercise appears to ‘prime’ effector T cells, thereby allow them to transmigrate
101 to the peripheral tissues that require enhanced immune surveillance following physical stress
102 (54). Compared to the resting condition, the percentage of circulating lymphocytes expressing
103 effector cytokines is lower following prolonged exercise (115), but it is unknown whether this
104 decline reflects impairment at the individual cell level or preferential movement of effector T
105 cells into peripheral tissues (e.g., lungs, gut). Recent evidence showing that exercise redeploys
106 T cells that are specific to latent herpesviruses such as cytomegalovirus (CMV) and Epstein–
107 Barr virus (EBV) (111, 112) suggests that this response may be a countermeasure against
108 stress-induced viral reactivation (107). Exercise may also mobilize ‘older’ functionally
109 exhausted/senescent lymphocytes to undergo apoptosis in the tissues and allow new
110 ‘recruits’ to take their place (106, 107).

111 Monocytes mobilized by exercise are likely to infiltrate skeletal muscle and differentiate
112 into tissue-resident macrophages that facilitate repair and regeneration, particular following
113 arduous bouts of exercise that cause significant skeletal muscle damage (85). Monocytes with
114 effector phenotypes are also preferentially redeployed after exercise. The CD14⁺/CD16⁺

115 'proinflammatory' monocytes are preferentially mobilized over their CD14⁺/CD16⁻
 116 counterparts (109). Monocyte expression of pathogen recognition receptors (e.g., toll-like
 117 receptors [TLRs]) tends to decrease in response to moderate-intensity exercise (109).
 118 Conversely, prolonged, intense exercise (60 km cycling time trial) increases TLR2 and TLR4
 119 expression on monocytes, which may indicate a heightened proinflammatory state (11). A
 120 recent study showed that acute exercise mobilizes angiogenic T cells, which may facilitate
 121 vascular remodeling during exercise recovery (53). Exercise is also known to mobilize
 122 hematopoietic stem cells, which may participate in skeletal muscle repair and regeneration
 123 after exercise (25, 49). It has been suggested that exercise may have a role as an adjuvant to
 124 mobilize stem cells in donors for hematopoietic stem cell transplantation (25).

125 In addition to cellular redeployment, the recovery phase of exercise, especially following
 126 very prolonged bouts of endurance-based exercise, is marked by striking alterations in the
 127 functional capacity of several blood leukocyte populations. Neutrophil bactericidal activity is
 128 greatly influenced by the intensity and duration of exercise. For example, after 1 h of cycling
 129 at 50% and 80% of $\dot{V}O_{2\max}$ increases and reduces neutrophil oxidative burst activity,
 130 respectively (94). During the early stages of recovery after exercise, neutrophil bactericidal
 131 activity continues to increase after 40 min to 1 h of moderate intensity exercise, whereas it
 132 remains impaired after exhaustive or prolonged exercise (86). NK cell cytotoxicity after
 133 exercise bouts of relatively short duration tends to remain unchanged on a per cell basis
 134 during recovery (81) but may decline after very prolonged bouts (33). T-cell proliferation in
 135 response to mitogen stimulation typically decreases both during and after exercise, regardless
 136 of exercise modality, intensity, or duration (126). Prolonged exercise may also reduce T-cell
 137 homing and migration (8), LPS-induced cytokine secretion by monocytes (113), and the

percentage of T cells producing effector cytokines in response to mitogen stimulation (115). Thus, the general trend during exercise recovery is that short bouts of moderate-intensity exercise have little effect (or might even enhance) cellular immune function, whereas prolonged bouts (>1.5 h) of heavy exertion appear to reduce the normal functioning of all major immune cell subtypes. These effects may leave athletes susceptible to illness during recovery from competition or heavy training (87).

144

145 **REPEATED EXERCISE BOUTS AND EXTENDED PERIODS OF INTENSE TRAINING**

The repeated bout paradigm (87) proposes that, compared with a single bout of exercise, repeated exercise bouts on the same day (27, 73, 96, 102) or over several days (43) cause different changes in circulating cell counts, lymphocyte proliferation, and NK cytotoxicity. Subsequent research has investigated changes in other immune responses to repeated exercise bouts on the same day with short versus long recovery and intensified training over weeks or months. Table 1 summarizes the evidence for changes in the immune system after repeated exercise bouts and days to months of intense training.

153 *One Versus Two Bouts of Exercise per Day*

Studies on the effects of one versus two bouts of exercise on a single day have included physically active (56-58, 63, 96, 102), highly trained (23, 88), or elite (12, 73, 98, 99) participants. The exercises involved cycling (12, 27, 56-58, 63, 96, 98, 99, 102), running (23, 88), or rowing (73). The duration and intensity ranged from <15 min at maximal intensity (27, 73) up to 2 h at medium–high intensity (i.e., 60–75% $\dot{V}O_{2\max}$) (56, 96). The recovery period between exercise bouts was most commonly 3–4 h but ranged from 45 min (102) to 12 h (88).

160 Compared with the initial bout of exercise, typically either show a greater relative change or
 161 a higher absolute value after a second bout of exercise for the following immune variables:
 162 total leukocyte count (57, 63, 99, 102), neutrophil count (23, 73, 96, 99, 102), oxidative burst
 163 (per neutrophil) (12), elastase release (per neutrophil) (57), CD4⁺ T-cell count (73, 99, 102),
 164 whole-blood IL-8 production (23), and NK cell activation (represented by CD69 expression)
 165 (99). Conversely, lymphocyte proliferation (96, 102) and whole-blood IL-6 production (23) are
 166 typically lower following a second bout compared with the first bout of exercise.

167 *Short Versus Long Recovery*

168 Three studies on the effects of recovery duration between two bouts of exercise included
 169 highly trained or elite athletes who cycled for 65 min at 75% $\dot{V}O_{2\max}$, twice each day, with
 170 either 3 or 6 h between the exercise bouts (12, 97, 98). A short recovery period (i.e., 3 h)
 171 induces either a greater relative increase or higher absolute values for neutrophil count (12,
 172 97), oxidative burst activity per neutrophil (12), and CD8⁺ T-cell and NK-cell counts (97) after
 173 the second bout of exercise. Exercise-induced changes in lymphocyte (97), monocyte, and
 174 eosinophil (12) counts; absolute oxidative burst activity (12); NK cytotoxicity (97); and plasma
 175 concentrations of IL-6 and IL-1ra (98) do not differ after a short versus long recovery period.

176 *Consecutive or Multiple Days of Exercise*

177 Studies detailing how the immune system responds to exercise repeated on consecutive
 178 days or every second day have included untrained or physically active participants (43, 116,
 179 118) or well-trained or elite athletes (60, 73, 77, 79). Exercise included 3 × 6 min maximal
 180 rowing, repeated twice over 2 d (73), 1–3 h cycling (77, 79, 118) or running (60) at 50–70%
 181 $\dot{V}O_{2\max}$ repeated over 3 d, every second day for 3 d (43), or daily for 7 d (116). Exercise-induced
 182 changes in plasma cytokine and elastase concentrations and cytokine mRNA expression in

183 leukocytes and muscle diminish over consecutive days (77, 118). By contrast, changes in total
184 leukocyte, neutrophil, and monocyte counts (73, 116, 118), lymphocyte proliferation (79),
185 neutrophil chemotaxis (116), leukocyte IL-1ra mRNA expression (77), plasma
186 myeloperoxidase concentration, and salivary IgA secretion rate (79) do not change over time.
187 Changes in lymphocyte subsets (43, 73), oxidative burst activity (79, 116, 118), salivary IgA
188 secretion rate (60, 77), and NK-cell count and cytotoxicity (73, 79) over consecutive days are
189 more variable.

190 *Immune Changes Associated with Overreaching and Overtraining*

191 Studies on short periods (2–4 weeks) of functional overreaching have reported decreases
192 in resting neutrophil degranulation (95), lymphocyte proliferation, and antibody production
193 (124). Neutrophil count, plasma cytokine concentrations, CD4:CD8 T-cell ratio, and salivary
194 IgA concentration are more variable (or do not change) in response to functional overreaching
195 (39, 95, 124). Athletes who exhibit signs of nonfunctional overreaching and/or frequent upper
196 respiratory illness present with lower salivary IgA concentration (26, 35, 60); lower cytokine
197 production by monocytes, neutrophils and dendritic cells (67); and a greater number of
198 activated (CD25⁺) lymphocytes (29). Changes in differential blood cell counts, lymphocyte
199 subsets, and NK-cell count following extended periods of intensified training are variable (29,
200 35, 61). Studies of athletes exhibiting the hallmarks of overtraining syndrome—including
201 illness—have not revealed any consistent or characteristic immune profile (30, 101).

202

203 SLEEP DISTURBANCE AND IMMUNE FUNCTION

204 Sleep disturbances influence immunity via activation of the hypothalamic–pituitary–
205 adrenal axis and the sympathetic nervous-system (46). Chronic sleep disturbance and
206 disruption to the normal circadian rhythm are associated with inflammation and
207 desynchronization of rhythmic immune variables. These responses likely contribute to
208 increased risk of infection, cardiovascular disease, and cancer in shift workers (21, 68).
209 Despite evidence that athletes experience poor sleep patterns compared with nonathletes
210 (16, 55), surprisingly little is known about how sleep disturbance influences the immune
211 responses to exercise. Compared with normal sleep, a disrupted night’s sleep appears to
212 prime the immune system and enhance immunosurveillance by stimulating total
213 lymphocytes, CD8⁺ T cells, NK cells, and $\gamma\delta$ T cells to leave the blood and migrate to potential
214 sites of infection during the early recovery period after exercise (45). By contrast, other
215 studies indicate that a night without sleep does not influence leukocyte trafficking, neutrophil
216 degranulation, or mucosal immunity at rest or after exercise (31, 93). Subtle immune changes
217 have been observed after a night without sleep, including a shift toward a T helper 2 cytokine
218 profile (46).

219 It is uncertain whether these subtle immune modifications with acute sleep loss are
220 clinically meaningful. When considering the potential effects of poor sleep on immunity in
221 athletes, it is important to distinguish between acute and chronic sleep disturbance. Chronic
222 sleep disturbance (12 nights, 50% sleep loss) increases the plasma inflammation markers C-
223 reactive protein and IL-6 (38). However, intervening daytime naps can counter this apparent
224 inflammatory response (103). Short sleep duration (<7 h/night for 7 d) decreases the response
225 to hepatitis B vaccination and the likelihood of clinical protection (90). Similarly, a night of

226 wakefulness after hepatitis A vaccination decreases the specific antibody response 2–4
 227 months later (52). People who experience poor quality sleep and/or regular sleep deprivation
 228 also have a 4–5-times greater risk of developing the common cold (16, 91). Continued
 229 research efforts should be directed toward monitoring and improving sleep in athletes and
 230 understanding the implications for immune health.

231

232 **NUTRITIONAL INTERVENTIONS FOR RESTORING IMMUNE FUNCTION AFTER EXERCISE**

233 Research over the last 30 years has investigated whether nutritional strategies counteract
 234 exercise-induced immunodepression and systemic inflammation (32, 125). A comprehensive
 235 review of the literature on these is beyond the scope of this mini-review; other more detailed
 236 reviews on this topic are available (e.g., (32, 125). Here, we focus on the most effective
 237 nutritional strategies—primarily carbohydrate ingestion—for restoring systemic immune
 238 function in the first few hours after exercise (9, 10, 19, 51, 65, 76, 78, 82, 83) and over
 239 consecutive days (7). We also assess whether the timing of nutritional interventions (i.e.,
 240 before, during or after exercise) influence their effectiveness (50, 59).

241 *Carbohydrate Supplementation before and/or during Exercise*

242 Carbohydrate supplementation during prolonged, intense exercise consistently attenuates
 243 exercise-induced increases in circulating cytokines (74, 125), and the re-distribution of
 244 neutrophils (74, 76, 78), monocytes (76, 82), natural killer cells (78), and lymphocytes (51).
 245 The immunomodulatory effects of carbohydrates arise from better maintenance of blood
 246 glucose concentrations and blunted release of stress hormones such as catecholamines and
 247 glucocorticoids during and after exercise (51, 76, 78, 82, 83). Although the systemic release

248 of IL-6 during exercise is related to muscle glycogen depletion (114), the precise mechanism
249 by which carbohydrate supplementation reduces systemic IL-6 release from contracting
250 muscle during exercise is not clear, because carbohydrate supplementation does not alter
251 muscle glycogen content (75).

252 In several studies, the immunomodulatory effects of carbohydrate supplementation
253 were observed to 'carry over' into the recovery period (i.e., ≥ 2 h post-exercise) (51, 76, 78,
254 82, 83). Nieman et al reported that carbohydrate supplementation during 2.5 h high-intensity
255 running reduces the number of neutrophils (immediately and 1.5 h post-exercise), monocytes
256 (immediately and 6 h post-exercise) and lymphocytes (immediately and 3 h post-exercise)
257 (76, 78). Extending carbohydrate ingestion to the post-exercise recovery period also reduces
258 neutrophil count (74), blood granulocyte and monocyte phagocytosis 6 h post-exercise (82).
259 Lancaster et al showed that carbohydrate consumption (30 and 60 g per hour) during 2.5 h
260 cycling minimized the suppression of CD4⁺ and CD8⁺ T lymphocyte that express and produce
261 IFN- γ during the 2 h following exercise (51).

262 Considering that exercise-induced responses of the adaptive immune system are relatively
263 slow (125), it is important to assess whether these effects are maintained over consecutive
264 days of exercise. Carbohydrate ingestion before, during and after two exercise bouts on two
265 consecutive days attenuated the decrease in antigen-stimulated proliferative lymphocyte
266 responses before exercise on the second day (7). Carbohydrate ingestion also enhanced
267 lymphocyte proliferative responses to mitogen stimulation post-exercise on the second day
268 (7). These findings suggest that carbohydrates may help to diminish potential cumulative
269 immunodepression over consecutive days of exercise.

270 The immunomodulatory effects of carbohydrate may depend on the timing of

271 carbohydrate intake. The ingestion of a glucose solution 15 min—but not 75 min—before 1 h
 272 high-intensity cycling prevented immunoendocrine perturbations (50). The lack of an effect
 273 of carbohydrates ingested 75 min pre-exercise was potentially associated with an insulin-
 274 induced decrease in the plasma glucose concentration prior to exercise, which in turn might
 275 have enhanced immunoendocrine responses (50). Carbohydrate ingestion during either the
 276 first or the second of two 90-min bouts of cycling on the same day better maintained plasma
 277 glucose and attenuated plasma stress hormone responses to the second bout (59). By
 278 contrast, carbohydrate ingestion during the 2 h recovery period between these exercise bouts
 279 had no such effects (59). These findings suggest beneficial effects of a timely carbohydrate
 280 supplementation (i.e., shortly before and/or during exercise) on immune responses to
 281 exercise. This may be particularly relevant with more prolonged and/or intense exercise
 282 protocols, and when the recovery duration between two consecutive exercise bouts is short.

283 Carbohydrate ingestion does not influence all aspects of the immune system. For example,
 284 carbohydrate supplementation does not alter the exercise-induced suppression of natural
 285 killer cell function (78) or salivary IgA secretion (18). Importantly, it remains unclear whether
 286 the immunomodulatory effects of carbohydrates have clinical relevance for resistance to
 287 illness or adaptation of the immune system to regular exercise stress (32, 125). Recent
 288 evidence indicates that carbohydrate supplementation during prolonged exercise blunts
 289 exercise-induced immune-endocrine perturbations, but does not prevent the suppression of
 290 in vivo immunity (22). More research is required to examine the effects of carbohydrates (or
 291 other nutritional strategies) on in vivo immune function in response to acute and chronic
 292 exercise.

293 *Dietary Carbohydrate Intake after Glycogen-Depletion*

294 Some studies have investigated the effects of dietary carbohydrate intake on immune
 295 responses to consecutive days of exercise intended to deplete muscle glycogen (9, 10, 34, 65).
 296 A higher carbohydrate intake consistently attenuated certain components of
 297 immunodepression well into the recovery period (i.e., ≥ 2 hours post-exercise) after the
 298 second exercise session (10, 34, 65). Athletic training often involves conditions of low
 299 carbohydrate availability e.g., due to abbreviated recovery periods and/or as part of ‘train
 300 low-compete high’ training regime (41, 42). These investigations therefore have particular
 301 practical implications. Compared with a higher carbohydrate intake (8 g/kg/d), very low
 302 carbohydrate intake (0.5 g/kg/d) leads to greater perturbation in leukocyte subsets during
 303 recovery from exercise (65). These effects may be related to sustained elevation of plasma
 304 cortisol concentration (65). Bishop et al observed that compared with a low-carbohydrate
 305 diet (1.1 g/kg/d), a high- (8.4 g/kg/d) for 3 d after glycogen-lowering cycling attenuated
 306 plasma cortisol and cytokine concentrations, circulating total leukocyte and neutrophil counts
 307 following subsequent exercise (10).

308 Consuming a high carbohydrate diet (8.5 g/kg/d) also reduces overreaching symptoms
 309 during 11 d of intense training, compared with moderate carbohydrate consumption (5.5
 310 g/kg/d) (1). The periodic implementation of ‘train-low’ strategies (e.g., by commencing
 311 training with low muscle glycogen stores) may further amplify metabolic adaptations in
 312 skeletal muscle (41, 42). When considering their dietary carbohydrate intake, athletes should
 313 aim to achieve a balance between minimizing immunodepression and maximizing metabolic
 314 adaptations in skeletal muscle. In view of the detrimental effects of low carbohydrate
 315 availability on the immune system, *chronic* carbohydrate restriction should be avoided during
 316 intense periods of training (32, 41). Additional research is warranted to better understand the
 317 effect of long-term training consisting of intermittent ‘train-low’ sessions on immune function

318 and susceptibility to illness.

319 *Dietary Protein Intake and Post-Exercise Protein Supplementation*

320 Recognizing the importance of protein for immunocompetence (15), there are benefits of
 321 post-exercise protein ingestion (18, 19, 69) or a diet high in protein (128) on immune
 322 responses to exercise. Based on previous results indicating that the exercise-induced
 323 lymphocyte trafficking was impaired during high-intensity training, Witard et al examined
 324 whether a high protein diet can restore these impaired immune responses (128). Consuming
 325 a high protein diet (3 g/kg/d) helped to minimize exercise-induced changes in lymphocyte
 326 distribution during a period of intense training (128). Interestingly, an energy- and
 327 carbohydrate-matched normal protein diet (1.5 g/kg/d) failed to provide the same benefit
 328 (128). The high-protein diet was also associated with fewer self-reported upper-respiratory
 329 illnesses (128). Another study demonstrated that protein and leucine supplementation for 1–
 330 3 h post-exercise during 6 d of high-intensity training enhanced neutrophil respiratory burst
 331 activity after the last exercise session (69). Consuming a carbohydrate-protein solution
 332 immediately—but not 1 h—after exercise prevents a decrease in neutrophil degranulation
 333 during the post-exercise recovery period (19).

334 Recent research has shown that the timing, distribution and amount of post-exercise
 335 protein intake modulate the blood and tissue availability of protein/amino acids and adaptive
 336 responses of skeletal muscle (3, 42). Notably, amino acid-sensitive mammalian target of
 337 rapamycin (mTOR) signaling is also a key mechanism underlying leukocyte trafficking (110).
 338 More studies are therefore needed to examine whether different post-exercise protein
 339 feeding patterns influence immune function during recovery from exercise.

340 *Antioxidants and Phytochemicals*

341 Except for carbohydrate supplementation, evidence for effective nutritional
342 countermeasures to exercise-induced immune alterations is limited (32, 125). Among other
343 types of nutritional supplements (e.g., probiotics and vitamin D (for reviews see (32, 125)),
344 antioxidants and phytochemicals such as quercetin have been studied for their potential
345 capacity to minimize immune perturbations—particularly during exercise recovery (75, 77,
346 79, 80). Some data point toward beneficial effects of quercetin supplementation on immune
347 health after intense exercise (77, 79). Other findings suggest an increased need for nutritional
348 antioxidants during the first 24 h of recovery from intense exercise lasting several hours such
349 as an Ironman triathlon (71). However, taken together, the present literature is not
350 sufficiently robust to recommend supplementation with phytochemicals or antioxidants to
351 prevent immune suppression and illness in athletes and exercising individuals. Athletes often
352 take high doses of antioxidant/phytochemical supplements in the belief that this will reduce
353 their risk of illness (47). However, high doses of antioxidant/phytochemical supplements can
354 interfere with training adaptations (42, 72). A natural diet rich in fruits, vegetables, whole
355 grains and nuts delivers antioxidants and phytochemicals in physiologically effective amounts
356 that are most likely sufficient to help maintain immune function following exercise and during
357 exercise training (32, 72, 125).

358

359 **OTHER STRATEGIES FOR RESTORING IMMUNE FUNCTION AFTER EXERCISE**

360 In addition to nutritional interventions, other research has examined the efficacy of
361 nonsteroidal anti-inflammatory drugs (NSAIDs) and various physical therapies for restoring
362 immune function after exercise. Some studies (13, 20, 70, 89, 120, 127) have shown effects
363 of NSAIDs, but other human studies have failed to demonstrate any effects of NSAIDs (122,

123), cryotherapy (44), compression garments (5, 37, 92), active recovery (2, 119), or other physical therapies (28) on immune responses during recovery from exercise (Fig. 2). Despite this lack of empirical evidence for the benefits of NSAIDs and physical therapies for restoring immune function after exercise, some of these treatments are associated with positive psychological outcomes and other effects not related to immune function after exercise (4, 17). Therefore, although the physiological effects of these physical treatments are not understood fully at the present time, they may confer some important benefits for athletes, which may involve the immune system—perhaps indirectly.

372

373 THEORETICAL AND PRACTICAL CONSIDERATIONS

374 The redeployment of effector lymphocytes from blood to peripheral tissues is seen as an integral part of the physiological stress response and the immune system's response to prepare the body for potential injury by mobilizing its 'troops' (cells) to increase immunosurveillance (24). Therefore, immune integrity during exercise recovery may be characterized by the host's ability to redeploy effector lymphocytes effectively to the peripheral tissues. Lymphocyte redeployment may be impaired following very prolonged bouts of exercise or in athletes who are overreaching. The redeployment of CD8⁺ T cells both during and after exercise is significantly reduced after 1 wk of high-intensity training compared with normal training (129). Specifically, the egress of CD8⁺ T cells was 1.4-fold higher after normal compared with high-intensity training.

384 Considering other evidence that stress-induced leukocyte redeployment is linked to poor clinical outcomes following surgery (100), immune cell redistribution and infection risk after

386 exercise warrants further investigation. High-volume exercise training may impair the
387 redeployment of viral-specific T cells and NK cells, thereby reducing antiviral ‘patrolling’
388 during exercise recovery (Fig. 1). Herpes viruses such as EBV and CMV are highly prevalent in
389 the population, and reactivation of these viruses from a latent state is indicative of systemic
390 immunodepression (107). EBV viral DNA is present in saliva from athletes after even short
391 periods of high-intensity training (36), but it is unknown whether this reflects training-induced
392 impairment in the trafficking of virus-fighting lymphocytes.

393 Cellular immune function in response to exercise is typically assessed in isolated cells ex
394 vivo, or at the cell population level in the blood compartment. This approach can make it
395 difficult to interpret changes in immune cell function after exercise because of the massive
396 alterations in the cellular composition of discrete leukocyte subtypes. On the one hand, it
397 seems intuitive to interpret lower immune cell function measured in blood during the early
398 stages of exercise recovery as indicative of immunodepression. On the other hand, it is equally
399 possible that, after exercise, the most functional immune cells (i.e., those with effector
400 phenotypes and high tissue-migrating potential) are redeployed to other areas of the body
401 where they are needed. If true, this suggests that systemic immunosurveillance may be
402 enhanced during exercise recovery, despite an apparent depressed profile in the blood
403 compartment.

404 It is difficult to determine the biological significance of exercise-induced changes in
405 immune cell function when assessed *in vitro*. This is because cell function is typically assessed
406 relative to the total cell population (e.g., percentage of T cells responding to mitogen
407 stimulation, number of target cells killed per NK-cell, oxidative burst activity per neutrophil,
408 etc.) without accounting for exercise-induced changes in the subset composition of these cell

409 populations. For example, the proportion of NK cells expressing the activating receptor,
410 NKG2C, is markedly elevated during exercise recovery (6). As a result, NK-cell cytotoxic activity
411 for the total NK-cell population increases markedly when an NKG2C-sensitive target cell is
412 used to assess NK-cell function (6). Conversely, when an NKG2C-insensitive target cell line
413 (K562) is used, NK-cell killing is not affected by exercise (6). Therefore, the proportional shifts
414 in the composition of cell subtypes should be considered when interpreting exercise-induced
415 changes in immune cell function at the total cell population level in vitro. Moreover,
416 assessment of in vitro immune cell function using venous blood samples does not account for
417 the complex interactions among immune cells and soluble factors within tissues (e.g., gut,
418 lungs, skin). However, some evidence suggests that reduced immune cell function in vitro may
419 coincide with changes in vivo and rates of illness (14, 40).

420 The validity of the original paradigm of cumulative immunodepression with repeated
421 bouts of exercise (87) is somewhat difficult to assess. Months of intense training increase the
422 incidence of illness in elite athletes (26, 30, 35). However, based on these studies, we can only
423 assume—but not assert—that increased incidence of illness results from an imbalance
424 between training and recovery. Research that has systematically manipulated the balance
425 between training and recovery has not identified any immune variables that are consistently
426 depressed as a result of insufficient recovery after exercise. However, with one exception
427 (79), these studies have not tracked the incidence of illness after repeated bouts of exercise.

428 Reduced salivary IgA concentration and secretion rate (amount of IgA secreted over a
429 fixed period) may predispose athletes to illness in the long term (26, 35). IgA binds micro-
430 organisms such as bacteria and viruses in the mucosa so that they can be destroyed by
431 immune cells. However, short-term changes in salivary IgA concentration and secretion rate

432 after repeated bouts of exercise are variable (56, 58, 60, 79). Salivary IgA concentration and
433 secretion rate may decrease incrementally over longer periods. The repeated exercise models
434 used in many of the studies described above induce only acute fatigue (64). Accordingly,
435 smaller exercise-induced changes in immune variables following repeated bouts of exercise
436 may actually represent positive adaptation of the immune system—as opposed to depression
437 of immunity that may lead to illness.

438 Pedersen et al (87) suggested that there is a critical threshold for exercise intensity and
439 duration that determines the risk of immunodepression after repeated bouts of exercise.
440 However, no studies have systematically determined the effects of repeated exercise bouts
441 of different intensity and duration. There are also no data on the effects of repeated bouts of
442 anaerobic or resistance/strength exercise, or a combination of different types of exercise on
443 the same day. The large gaps in Table 1 show that much remains to be learned about the
444 effects of repeated bouts of exercise on the immune system.

445 Among various nutritional interventions that have been studied to counteract
446 immunodepression during exercise recovery, carbohydrate supplementation has proven the
447 most effective. A balanced and well-diversified diet that meets the energy demands in
448 athletes and exercising individuals is certainly a key component to maintain immune function
449 in response to strenuous exercise and intense periods of training. Additional research is
450 warranted to investigate how the timing and pattern in the ingestion of nutrients—
451 particularly carbohydrates and protein/amino acids—influence recovery of the immune
452 system after exercise.

453 Sleep disturbances can depress immunity, increase inflammation, and promote adverse
454 health outcomes in the general population. However, the limited data available on how sleep

455 disturbances influence immune responses to exercise are inconsistent. Physical treatments
456 that are used after exercise (e.g., hydrotherapy and massage) may enhance the athlete's
457 sense of well-being, and should be considered as adjunct therapies for maintaining immune
458 health.

459 **ACKNOWLEDGMENTS**

460 Jonathan Peake is supported by funding from the Centre of Excellence for Applied Sport
461 Science Research at the Queensland Academy of Sport, Brisbane, Australia. We acknowledge
462 copyediting by Laurel T Mackinnon, PhD, FACSM, ELS.

463 REFERENCES

464 1. **Achten J, Halson SL, Moseley L, Rayson MP, Casey A, and Jeukendrup AE.** Higher dietary
465 carbohydrate content during intensified running training results in better maintenance of
466 performance and mood state. *J Appl Physiol* (1985) 96: 1331-1340, 2004.

467 2. **Andersson H, Bohn SK, Raastad T, Paulsen G, Blomhoff R, and Kadi F.** Differences in the
468 inflammatory plasma cytokine response following two elite female soccer games separated by a 72-h
469 recovery. *Scand J Med Sci Sports* 20: 740-747, 2010.

470 3. **Areta JL, Burke LM, Ross ML, Camera DM, West DW, Broad EM, Jeacocke NA, Moore DR,**
471 **Stellingwerff T, Phillips SM, Hawley JA, and Coffey VG.** Timing and distribution of protein ingestion
472 during prolonged recovery from resistance exercise alters myofibrillar proteinsynthesis. *J Physiol* 591:
473 2319-2331, 2013.

474 4. **Arroyo-Morales M, Olea N, Ruiz C, del Castillo Jde D, Martinez M, Lorenzo C, and Diaz-Rodriguez**
475 **L.** Massage after exercise responses of immunologic and endocrine markers: a randomized single-blind
476 placebo-controlled study. *J Strength Cond Res* 23: 638-644, 2009.

477 5. **Bieuzen F, Brisswalter J, Easthope C, Vercruyssen F, Bernard T, and Hausswirth C.** Effect of
478 wearing compression stockings on recovery after mild exercise-induced muscle damage. *Int J Sports*
479 *Physiol Perform* 9: 256-264, 2014.

480 6. **Bigley AB, Rezvani K, Chew C, Sekine T, Pistillo M, Crucian B, Bollard CM, and Simpson RJ.** Acute
481 exercise preferentially redeploys NK-cells with a highly-differentiated phenotype and augments
482 cytotoxicity against lymphoma and multiple myeloma target cells. *Brain Behav Immun* 39: 160-171,
483 2014.

484 7. **Bishop NC, Walker GJ, Bowley LA, Evans KF, Molyneux K, Wallace FA, and Smith AC.** Lymphocyte
485 responses to influenza and tetanus toxoid in vitro following intensive exercise and carbohydrate
486 ingestion on consecutive days. *J Appl Physiol* (1985) 99: 1327-1335, 2005.

487 8. **Bishop NC, Walker GJ, Gleeson M, Wallace FA, and Hewitt CR.** Human T lymphocyte migration
488 towards the supernatants of human rhinovirus infected airway epithelial cells: influence of exercise
489 and carbohydrate intake. *Exerc Immunol Rev* 15: 127-144, 2009.

490 9. **Bishop NC, Walsh NP, Haines DL, Richards EE, and Gleeson M.** Pre-exercise carbohydrate status
491 and immune responses to prolonged cycling: I. Effect on neutrophil degranulation. *Int J Sport Nutr*
492 *Exerc Metab* 11: 490-502, 2001.

493 10. **Bishop NC, Walsh NP, Haines DL, Richards EE, and Gleeson M.** Pre-exercise carbohydrate status
494 and immune responses to prolonged cycling: II. Effect on plasma cytokine concentration. *Int J Sport*
495 *Nutr Exerc Metab* 11: 503-512, 2001.

496 11. **Booth S, Florida-James GD, McFarlin BK, Spielmann G, O'Connor DP, and Simpson RJ.** The impact
497 of acute strenuous exercise on TLR2, TLR4 and HLA-DR expression on human blood monocytes induced
498 by autologous serum. *Eur J Appl Physiol* 110: 1259-1268, 2010.

499 12. **Boyum A, Ronsen O, Tennfjord VA, Tollefsen S, Haugen AH, Opstad PK, and Bahr R.**
500 Chemiluminescence response of granulocytes from elite athletes during recovery from one or two
501 intense bouts of exercise. *Eur J Appl Physiol* 88: 20-28, 2002.

502 13. **Broadbent S, Rousseau JJ, Thorp RM, Choate SL, Jackson FS, and Rowlands DS.** Vibration therapy
503 reduces plasma IL6 and muscle soreness after downhill running. *Br J Sports Med* 44: 888-894, 2010.

504 14. **Bruunsgaard H, Hartkopp A, Mohr T, Konradsen H, Heron I, Mordhorst CH, and Pedersen BK.** In
505 vivo cell-mediated immunity and vaccination response following prolonged, intense exercise. *Med Sci*
506 *Sports Exerc* 29: 1176-1181, 1997.

507 15. **Calder PC, and Jackson AA.** Undernutrition, infection and immune function. *Nutr Res Rev* 13: 3-
508 29, 2000.

509 16. **Cohen S, Doyle WJ, Alper CM, Janicki-Deverts D, and Turner RB.** Sleep habits and susceptibility to
510 the common cold. *Arch Intern Med* 169: 62-67, 2009.

- 511 17. **Cortis C, Tessitore A, D'Artibale E, Meeusen R, and Capranica L.** Effects of post-exercise recovery
512 interventions on physiological, psychological, and performance parameters. *Int J Sports Med* 31: 327-
513 335, 2010.
- 514 18. **Costa RJ, Fortes MB, Richardson K, Bilzon JL, and Walsh NP.** The effects of postexercise feeding
515 on saliva antimicrobial proteins. *Int J Sport Nutr Exerc Metab* 22: 184-191, 2012.
- 516 19. **Costa RJ, Oliver SJ, Laing SJ, Waiters R, Bilzon JL, and Walsh NP.** Influence of timing of postexercise
517 carbohydrate-protein ingestion on selected immune indices. *Int J Sport Nutr Exerc Metab* 19: 366-384,
518 2009.
- 519 20. **Crane JD, Ogborn DI, Cupido C, Melov S, Hubbard A, Bourgeois JM, and Tarnopolsky MA.**
520 Massage therapy attenuates inflammatory signaling after exercise-induced muscle damage. *Sci Transl*
521 *Med* 4: 119ra113, 2012.
- 522 21. **Cuesta M, Boudreau P, Dubeau-Laramée G, Cermakian N, and Boivin DB.** Simulated night shift
523 disrupts circadian rhythms of immune functions in humans. *J Immunol* 196: 2466-2475, 2016.
- 524 22. **Davison G, Kehaya C, Diment BC, and Walsh NP.** Carbohydrate supplementation does not blunt
525 the prolonged exercise-induced reduction of in vivo immunity. *Eur J Nutr* 55: 1583-1593, 2016.
- 526 23. **Degerstrom J, and Osterud B.** Increased inflammatory response of blood cells to repeated bout of
527 endurance exercise. *Med Sci Sports Exerc* 38: 1297-1303, 2006.
- 528 24. **Dhabhar FS.** Enhancing versus Suppressive Effects of Stress on Immune Function: Implications for
529 Immunoprotection versus Immunopathology. *Allergy Asthma Clin Immunol* 4: 2-11, 2008.
- 530 25. **Emmons R, Niemiro GM, Owolabi O, and De Lisio M.** Acute exercise mobilizes hematopoietic stem
531 and progenitor cells and alters the mesenchymal stromal cell secretome. *J Appl Physiol* 120: 624-632,
532 2016.
- 533 26. **Fahlman MM, and Engels HJ.** Mucosal IgA and URTI in American college football players: a year
534 longitudinal study. *Med Sci Sports Exerc* 37: 374-380, 2005.
- 535 27. **Field CJ, Gougeon R, and Marliss EB.** Circulating mononuclear cell numbers and function during
536 intense exercise and recovery. *J Appl Physiol* 71: 1089-1097, 1991.
- 537 28. **Finberg M, Braham R, Goodman C, Gregory P, and Peeling P.** Effects of electrostimulation therapy
538 on recovery from acute team-sport activity. *Int J Sports Physiol Perform* 8: 293-299, 2013.
- 539 29. **Fry RW, Grove JR, Morton AR, Zeroni PM, Gaudieri S, and Keast D.** Psychological and
540 immunological correlates of acute overtraining. *Br J Sports Med* 28: 241-246, 1994.
- 541 30. **Gabriel HH, Urhausen A, Valet G, Heidelberg U, and Kindermann W.** Overtraining and immune
542 system: a prospective longitudinal study in endurance athletes. *Med Sci Sports Exerc* 30: 1151-1157,
543 1998.
- 544 31. **Gillum TL, Kuennen MR, Castillo MN, Williams NL, and Jordan-Patterson AT.** Exercise, but not
545 acute sleep loss, increases salivary antimicrobial protein secretion. *J Strength Cond Res* 29: 1359-1366,
546 2015.
- 547 32. **Gleeson M.** Immunological aspects of sport nutrition. *Immunol Cell Biol* 94: 117-123, 2016.
- 548 33. **Gleeson M, and Bishop NC.** The T cell and NK cell immune response to exercise. *Ann Transplant*
549 10: 43-48, 2005.
- 550 34. **Gleeson M, Blannin AK, Walsh NP, Bishop NC, and Clark AM.** Effect of low- and high-carbohydrate
551 diets on the plasma glutamine and circulating leukocyte responses to exercise. *Int J Sport Nutr* 8: 49-
552 59, 1998.
- 553 35. **Gleeson M, McDonald WA, Cripps AW, Pyne DB, Clancy RL, and Fricker PA.** The effect on
554 immunity of long-term intensive training in elite swimmers. *Clin Exp Immunol* 102: 210-216, 1995.
- 555 36. **Gleeson M, Pyne DB, Austin JP, Lynn Francis J, Clancy RL, McDonald WA, and Fricker PA.** Epstein-
556 Barr virus reactivation and upper-respiratory illness in elite swimmers. *Med Sci Sports Exerc* 34: 411-
557 417, 2002.
- 558 37. **Goto K, and Morishima T.** Compression garment promotes muscular strength recovery after
559 resistance exercise. *Med Sci Sports Exerc* 46: 2265-2270, 2014.

- 560 38. **Haack M, Sanchez E, and Mullington JM.** Elevated inflammatory markers in response to prolonged
561 sleep restriction are associated with increased pain experience in healthy volunteers. *Sleep* 30: 1145-
562 1152, 2007.
- 563 39. **Halsen SL, Lancaster GI, Jeukendrup AE, and Gleeson M.** Immunological responses to
564 overreaching in cyclists. *Med Sci Sports Exerc* 35: 854-861, 2003.
- 565 40. **Harper Smith AD, Coakley SL, Ward MD, Macfarlane AW, Friedmann PS, and Walsh NP.** Exercise-
566 induced stress inhibits both the induction and elicitation phases of in vivo T-cell-mediated immune
567 responses in humans. *Brain Behav Immun* 25: 1136-1142, 2011.
- 568 41. **Hawley JA, and Burke LM.** Carbohydrate availability and training adaptation: effects on cell
569 metabolism. *Exerc Sport Sci Rev* 38: 152-160, 2010.
- 570 42. **Hawley JA, Burke LM, Phillips SM, and Spriet LL.** Nutritional modulation of training-induced
571 skeletal muscle adaptations. *J Appl Physiol* (1985) 110: 834-845, 2011.
- 572 43. **Hoffman-Goetz L, Simpson JR, Cipp N, Arumugam Y, and Houston ME.** Lymphocyte subset
573 responses to repeated submaximal exercise in men. *J Appl Physiol* 68: 1069-1074, 1990.
- 574 44. **Hohenauer E, Taeymans J, Baeyens JP, Clarys P, and Clijsen R.** The effect of post-exercise
575 cryotherapy on recovery characteristics: a systematic review and meta-analysis. *PLoS One* 10:
576 e0139028, 2015.
- 577 45. **Ingram LA, Simpson RJ, Malone E, and Florida-James GD.** Sleep disruption and its effect on
578 lymphocyte redeployment following an acute bout of exercise. *Brain Behav Immun* 47: 100-108, 2015.
- 579 46. **Irwin MR.** Why sleep is important for health: a psychoneuroimmunology perspective. *Annu Rev*
580 *Psychol* 66: 143-172, 2015.
- 581 47. **Knez WL, and Peake JM.** The prevalence of vitamin supplementation in ultraendurance triathletes.
582 *Int J Sport Nutr Exerc Metab* 20: 507-514, 2010.
- 583 48. **Kruger K, Lechtermann A, Fobker M, Volker K, and Mooren FC.** Exercise-induced redistribution of
584 T lymphocytes is regulated by adrenergic mechanisms. *Brain Behav Immun* 22: 324-338, 2008.
- 585 49. **Kruger K, Pilat C, Schild M, Lindner N, Frech T, Muders K, and Mooren FC.** Progenitor cell
586 mobilization after exercise is related to systemic levels of G-CSF and muscle damage. *Scand J Med Sci*
587 *Sports* 25: e283-291, 2015.
- 588 50. **Lancaster GI, Jentjens RL, Moseley L, Jeukendrup AE, and Gleeson M.** Effect of pre-exercise
589 carbohydrate ingestion on plasma cytokine, stress hormone, and neutrophil degranulation responses
590 to continuous, high-intensity exercise. *Int J Sport Nutr Exerc Metab* 13: 436-453, 2003.
- 591 51. **Lancaster GI, Khan Q, Drysdale PT, Wallace F, Jeukendrup AE, Drayson MT, and Gleeson M.** Effect
592 of prolonged exercise and carbohydrate ingestion on type 1 and type 2 T lymphocyte distribution and
593 intracellular cytokine production in humans. *J Appl Physiol* (1985) 98: 565-571, 2005.
- 594 52. **Lange T, Dimitrov S, Bollinger T, Diekelmann S, and Born J.** Sleep after vaccination boosts
595 immunological memory. *J Immunol* 187: 283-290, 2011.
- 596 53. **Lansford KA, Shill DD, Dicks AB, Marshburn MP, Southern WM, and Jenkins NT.** Effect of acute
597 exercise on circulating angiogenic cell and microparticle populations. *Exp Physiol* 101: 155-167, 2016.
- 598 54. **LaVoy EC, Bosch JA, Lowder TW, and Simpson RJ.** Acute aerobic exercise in humans increases
599 cytokine expression in CD27(-) but not CD27(+) CD8(+) T-cells. *Brain Behav Immun* 27: 54-62, 2013.
- 600 55. **Leeder J, Glaister M, Pizzoferro K, Dawson J, and Pedlar C.** Sleep duration and quality in elite
601 athletes measured using wristwatch actigraphy. *J Sports Sci* 30: 541-545, 2012.
- 602 56. **Li TL, and Gleeson M.** The effect of single and repeated bouts of prolonged cycling and circadian
603 variation on saliva flow rate, immunoglobulin A and alpha-amylase responses. *J Sports Sci* 22: 1015-
604 1024, 2004.
- 605 57. **Li TL, and Gleeson M.** The effect of single and repeated bouts of prolonged cycling on leukocyte
606 redistribution, neutrophil degranulation, IL-6, and plasma stress hormone responses. *Int J Sport Nutr*
607 *Exerc Metab* 14: 501-516, 2004.
- 608 58. **Li TL, and Gleeson M.** The effects of carbohydrate supplementation during repeated bouts of
609 prolonged exercise on saliva flow rate and immunoglobulin A. *J Sports Sci* 23: 713-722, 2005.

- 610 59. **Li TL, and Gleeson M.** The effects of carbohydrate supplementation during the second of two
611 prolonged cycling bouts on immunoendocrine responses. *Eur J Appl Physiol* 95: 391-399, 2005.
- 612 60. **Mackinnon LT, and Hooper S.** Mucosal (secretory) immune system responses to exercise of
613 varying intensity and during overtraining. *Int J Sports Med* 15 Suppl 3: S179-183, 1994.
- 614 61. **Mackinnon LT, Hooper SL, Jones S, Gordon RD, and Bachmann AW.** Hormonal, immunological,
615 and hematological responses to intensified training in elite swimmers. *Med Sci Sports Exerc* 29: 1637-
616 1645, 1997.
- 617 62. **Mars M, Govender S, Weston A, Naicker V, and Chuturgoon A.** High intensity exercise: a cause of
618 lymphocyte apoptosis? *Biochem Biophys Res Commun* 249: 366-370, 1998.
- 619 63. **McFarlin B, Mitchell JB, McFarlin MA, and Steinhoff GM.** Repeated endurance exercise affects
620 leukocyte number but not NK cell activity. *Med Sci Sports Exerc* 35: 1130-1138, 2003.
- 621 64. **Meeusen R, Duclos M, Foster C, Fry A, Gleeson M, Nieman D, Raglin J, Rietjens G, Steinacker J,
622 and Urhausen A.** Prevention, diagnosis, and treatment of the overtraining syndrome: joint consensus
623 statement of the European College of Sport Science and the American College of Sports Medicine.
624 *Med Sci Sports Exerc* 45: 186-205, 2013.
- 625 65. **Mitchell JB, Pizza FX, Paquet A, Davis BJ, Forrest MB, and Braun WA.** Influence of carbohydrate
626 status on immune responses before and after endurance exercise. *J Appl Physiol* (1985) 84: 1917-1925,
627 1998.
- 628 66. **Mooren FC, Bloming D, Lechtermann A, Lerch MM, and Volker K.** Lymphocyte apoptosis after
629 exhaustive and moderate exercise. *J Appl Physiol* 93: 147-153, 2002.
- 630 67. **Morgado JM, Rama L, Silva I, de Jesus Inacio M, Henriques A, Laranjeira P, Pedreiro S, Rosado F,
631 Alves F, Gleeson M, Pais ML, Paiva A, and Teixeira AM.** Cytokine production by monocytes,
632 neutrophils, and dendritic cells is hampered by long-term intensive training in elite swimmers. *Eur J*
633 *Appl Physiol* 112: 471-482, 2012.
- 634 68. **Mullington JM, Simpson NS, Meier-Ewert HK, and Haack M.** Sleep loss and inflammation. *Best*
635 *Pract Res Clin Endocrinol Metab* 24: 775-784, 2010.
- 636 69. **Nelson AR, Jackson L, Clarke J, Stellingwerff T, Broadbent S, and Rowlands DS.** Effect of post-
637 exercise protein-leucine feeding on neutrophil function, immunomodulatory plasma metabolites and
638 cortisol during a 6-day block of intense cycling. *Eur J Appl Physiol* 113: 2211-2222, 2013.
- 639 70. **Nemet D, Meckel Y, Bar-Sela S, Zaldivar F, Cooper DM, and Eliakim A.** Effect of local cold-pack
640 application on systemic anabolic and inflammatory response to sprint-interval training: a prospective
641 comparative trial. *Eur J Appl Physiol* 107: 411-417, 2009.
- 642 71. **Neubauer O, Reichhold S, Nics L, Hoelzl C, Valentini J, Stadlmayr B, Knasmüller S, and Wagner**
643 **KH.** Antioxidant responses to an acute ultra-endurance exercise: impact on DNA stability and
644 indications for an increased need for nutritive antioxidants in the early recovery phase. *Br J Nutr* 104:
645 1129-1138, 2010.
- 646 72. **Neubauer O, and Yfanti C.** Antioxidants in Athlete's Basic Nutrition: Considerations towards a
647 Guideline for the Intake of Vitamin C and Vitamin E. In: *Antioxidants in Sport Nutrition*, edited by
648 Lamprecht M. Boca Raton (FL): 2015.
- 649 73. **Nielsen HB, Secher NH, Christensen NJ, and Pedersen BK.** Lymphocytes and NK cell activity during
650 repeated bouts of maximal exercise. *Am J Physiol* 271: R222-227, 1996.
- 651 74. **Nieman DC.** Influence of carbohydrate on the immune response to intensive, prolonged exercise.
652 *Exerc Immunol Rev* 4: 64-76, 1998.
- 653 75. **Nieman DC, Davis JM, Henson DA, Walberg-Rankin J, Shute M, Dumke CL, Utter AC, Vinci DM,
654 Carson JA, Brown A, Lee WJ, McNulty SR, and McNulty LS.** Carbohydrate ingestion influences
655 skeletal muscle cytokine mRNA and plasma cytokine levels after a 3-h run. *J Appl Physiol* 94: 1917-
656 1925, 2003.
- 657 76. **Nieman DC, Fagoaga OR, Butterworth DE, Warren BJ, Utter A, Davis JM, Henson DA, and**
658 **Nehlsen-Cannarella SL.** Carbohydrate supplementation affects blood granulocyte and monocyte
659 trafficking but not function after 2.5 h of running. *Am J Clin Nutr* 66: 153-159, 1997.

- 660 77. Nieman DC, Henson DA, Davis JM, Angela Murphy E, Jenkins DP, Gross SJ, Carmichael MD,
 661 Quindry JC, Dumke CL, Utter AC, McAnulty SR, McAnulty LS, Triplett NT, and Mayer EP. Quercetin's
 662 influence on exercise-induced changes in plasma cytokines and muscle and leukocyte cytokine mRNA.
 663 *J Appl Physiol* 103: 1728-1735, 2007.
- 664 78. Nieman DC, Henson DA, Garner EB, Butterworth DE, Warren BJ, Utter A, Davis JM, Fagoaga OR,
 665 and Nehlsen-Cannarella SL. Carbohydrate affects natural killer cell redistribution but not activity after
 666 running. *Med Sci Sports Exerc* 29: 1318-1324, 1997.
- 667 79. Nieman DC, Henson DA, Gross SJ, Jenkins DP, Davis JM, Murphy EA, Carmichael MD, Dumke CL,
 668 Utter AC, McAnulty SR, McAnulty LS, and Mayer EP. Quercetin reduces illness but not immune
 669 perturbations after intensive exercise. *Med Sci Sports Exerc* 39: 1561-1569, 2007.
- 670 80. Nieman DC, Henson DA, McAnulty SR, McAnulty LS, Morrow JD, Ahmed A, and Heward CB.
 671 Vitamin E and immunity after the Kona Triathlon World Championship. *Med Sci Sports Exerc* 36: 1328-
 672 1335, 2004.
- 673 81. Nieman DC, Miller AR, Henson DA, Warren BJ, Gusewitch G, Johnson RL, Davis JM, Butterworth
 674 DE, and Nehlsen-Cannarella SL. Effects of high- vs moderate-intensity exercise on natural killer cell
 675 activity. *Med Sci Sports Exerc* 25: 1126-1134, 1993.
- 676 82. Nieman DC, Nehlsen-Cannarella SL, Fagoaga OR, Henson DA, Utter A, Davis JM, Williams F, and
 677 Butterworth DE. Effects of mode and carbohydrate on the granulocyte and monocyte response to
 678 intensive, prolonged exercise. *J Appl Physiol* (1985) 84: 1252-1259, 1998.
- 679 83. Nieman DC, Nehlsen-Cannarella SL, Fagoaga OR, Henson DA, Utter A, Davis JM, Williams F, and
 680 Butterworth DE. Influence of mode and carbohydrate on the cytokine response to heavy exertion.
 681 *Med Sci Sports Exerc* 30: 671-678, 1998.
- 682 84. Peake J. Interrelations between acute and chronic stress and the immune and endocrine systems.
 683 In: *Endocrinology of physical activity and sport*, edited by Constantini N, and Hackney A. New York:
 684 Springer, 2013, p. 258-280.
- 685 85. Peake J, Nosaka K, and Suzuki K. Characterization of inflammatory responses to eccentric exercise
 686 in humans. *Exerc Immunol Rev* 11: 64-85, 2005.
- 687 86. Peake JM. Exercise-induced alterations in neutrophil degranulation and respiratory burst activity:
 688 possible mechanisms of action. *Exerc Immunol Rev* 8: 49-100, 2002.
- 689 87. Pedersen BK, Rohde T, and Ostrowski K. Recovery of the immune system after exercise. *Acta*
 690 *Physiol Scand* 162: 325-332, 1998.
- 691 88. Peeling P, Dawson B, Goodman C, Landers G, Wiegerinck ET, Swinkels DW, and Trinder D.
 692 Cumulative effects of consecutive running sessions on hemolysis, inflammation and hepcidin activity.
 693 *Eur J Appl Physiol* 106: 51-59, 2009.
- 694 89. Pournot H, Bieuzen F, Louis J, Mounier R, Fillard JR, Barbiche E, and Hausswirth C. Time-course
 695 of changes in inflammatory response after whole-body cryotherapy multi exposures following severe
 696 exercise. *PLoS One* 6: e22748, 2011.
- 697 90. Prather AA, Hall M, Fury JM, Ross DC, Muldoon MF, Cohen S, and Marsland AL. Sleep and
 698 antibody response to hepatitis B vaccination. *Sleep* 35: 1063-1069, 2012.
- 699 91. Prather AA, Janicki-Deverts D, Hall MH, and Cohen S. Behaviorally assessed sleep and
 700 susceptibility to the common cold. *Sleep* 38: 1353-1359, 2015.
- 701 92. Pruscino CL, Halson S, and Hargreaves M. Effects of compression garments on recovery following
 702 intermittent exercise. *Eur J Appl Physiol* 113: 1585-1596, 2013.
- 703 93. Ricardo JS, Cartner L, Oliver SJ, Laing SJ, Walters R, Bilzon JL, and Walsh NP. No effect of a 30-h
 704 period of sleep deprivation on leukocyte trafficking, neutrophil degranulation and saliva IgA responses
 705 to exercise. *Eur J Appl Physiol* 105: 499-504, 2009.
- 706 94. Robson PJ, Blannin AK, Walsh NP, Castell LM, and Gleeson M. Effects of exercise intensity,
 707 duration and recovery on in vitro neutrophil function in male athletes. *Int J Sports Med* 20: 128-135,
 708 1999.
- 709 95. Robson-Ansley PJ, Blannin A, and Gleeson M. Elevated plasma interleukin-6 levels in trained male
 710 triathletes following an acute period of intense interval training. *Eur J Appl Physiol* 99: 353-360, 2007.

711 96. **Rohde T, MacLean DA, and Pedersen BK.** Effect of glutamine supplementation on changes in the
712 immune system induced by repeated exercise. *Med Sci Sports Exerc* 30: 856-862, 1998.

713 97. **Ronsen O, Kjeldsen-Kragh J, Haug E, Bahr R, and Pedersen BK.** Recovery time affects
714 immunoendocrine responses to a second bout of endurance exercise. *Am J Physiol Cell Physiol* 283:
715 C1612-1620, 2002.

716 98. **Ronsen O, Lea T, Bahr R, and Pedersen BK.** Enhanced plasma IL-6 and IL-1ra responses to repeated
717 vs. single bouts of prolonged cycling in elite athletes. *J Appl Physiol* 92: 2547-2553, 2002.

718 99. **Ronsen O, Pedersen BK, Oritsland TR, Bahr R, and Kjeldsen-Kragh J.** Leukocyte counts and
719 lymphocyte responsiveness associated with repeated bouts of strenuous endurance exercise. *J Appl*
720 *Physiol* 91: 425-434, 2001.

721 100. **Rosenberger PH, Ickovics JR, Epel E, Nadler E, Jokl P, Fulkerson JP, Tillie JM, and Dhabhar FS.**
722 Surgical stress-induced immune cell redistribution profiles predict short-term and long-term
723 postsurgical recovery. A prospective study. *J Bone Joint Surg Am* 91: 2783-2794, 2009.

724 101. **Rowbottom DG, Keast D, Goodman C, and Morton AR.** The haematological, biochemical and
725 immunological profile of athletes suffering from the overtraining syndrome. *Eur J Appl Physiol Occup*
726 *Physiol* 70: 502-509, 1995.

727 102. **Severs Y, Brenner I, Shek PN, and Shephard RJ.** Effects of heat and intermittent exercise on
728 leukocyte and sub-population cell counts. *Eur J Appl Physiol Occup Physiol* 74: 234-245, 1996.

729 103. **Shearer WT, Reuben JM, Mullington JM, Price NJ, Lee BN, Smith EO, Szuba MP, Van Dongen**
730 **HP, and Dinges DF.** Soluble TNF-alpha receptor 1 and IL-6 plasma levels in humans subjected to the
731 sleep deprivation model of spaceflight. *J Allergy Clin Immunol* 107: 165-170, 2001.

732 104. **Shek PN, Sabiston BH, Buguet A, and Radomski MW.** Strenuous exercise and immunological
733 changes: a multiple-time-point analysis of leukocyte subsets, CD4/CD8 ratio, immunoglobulin
734 production and NK cell response. *Int J Sports Med* 16: 466-474, 1995.

735 105. **Simpson R, Florida-James G, Whyte G, Black J, Ross J, and Guy K.** Apoptosis does not
736 contribute to the blood lymphocytopenia observed after intensive and downhill treadmill running in
737 humans. *Res Sports Med* 15: 157-174, 2007.

738 106. **Simpson RJ.** Aging, persistent viral infections, and immunosenescence: can exercise "make
739 space"? *Exerc Sport Sci Rev* 39: 23-33, 2011.

740 107. **Simpson RJ, Bigley AB, Spielmann G, LaVoy EC, Kunz H, and Bollard CM.** Human
741 cytomegalovirus infection and the immune response to exercise. *Exerc Immunol Rev* 22: 8-27, 2016.

742 108. **Simpson RJ, Florida-James GD, Whyte GP, and Guy K.** The effects of intensive, moderate and
743 downhill treadmill running on human blood lymphocytes expressing the adhesion/activation
744 molecules CD54 (ICAM-1), CD18 (beta2 integrin) and CD53. *Eur J Appl Physiol* 97: 109-121, 2006.

745 109. **Simpson RJ, McFarlin BK, McSporran C, Spielmann G, o Hartaigh B, and Guy K.** Toll-like
746 receptor expression on classic and pro-inflammatory blood monocytes after acute exercise in humans.
747 *Brain Behav Immun* 23: 232-239, 2009.


748 110. **Sinclair LV, Finlay D, Feijoo C, Cornish GH, Gray A, Ager A, Okkenhaug K, Hagenbeek TJ, Spits**
749 **H, and Cantrell DA.** Phosphatidylinositol-3-OH kinase and nutrient-sensing mTOR pathways control T
750 lymphocyte trafficking. *Nat Immunol* 9: 513-521, 2008.

751 111. **Spielmann G, Bollard CM, Bigley AB, Hanley PJ, Blaney JW, LaVoy EC, Pircher H, and Simpson**
752 **RJ.** The effects of age and latent cytomegalovirus infection on the redeployment of CD8+ T cell subsets
753 in response to acute exercise in humans. *Brain Behav Immun* 39: 142-151, 2014.

754 112. **Spielmann G, Bollard CM, Kunz H, Hanley PJ, and Simpson RJ.** A single exercise bout enhances
755 the manufacture of viral-specific T-cells from healthy donors: implications for allogeneic adoptive
756 transfer immunotherapy. *Sci Rep* 6: 25852, 2016.

757 113. **Starkie RL, Angus DJ, Rolland J, Hargreaves M, and Febbraio MA.** Effect of prolonged,
758 submaximal exercise and carbohydrate ingestion on monocyte intracellular cytokine production in
759 humans. *J Physiol* 528: 647-655, 2000.

114. **Steensberg A, Febbraio MA, Osada T, Schjerling P, van Hall G, Saltin B, and Pedersen BK.** Interleukin-6 production in contracting human skeletal muscle is influenced by pre-exercise muscle glycogen content. *J Physiol* 537: 633-639, 2001.
115. **Steensberg A, Toft AD, Bruunsgaard H, Sandmand M, Halkjaer-Kristensen J, and Pedersen BK.** Strenuous exercise decreases the percentage of type 1 T cells in the circulation. *J Appl Physiol* 91: 1708-1712, 2001.
116. **Suzuki K, Naganuma S, Totsuka M, Suzuki KJ, Mochizuki M, Shiraishi M, Nakaji S, and Sugawara K.** Effects of exhaustive endurance exercise and its one-week daily repetition on neutrophil count and functional status in untrained men. *Int J Sports Med* 17: 205-212, 1996.
117. **Suzuki K, Nakaji S, Yamada M, Liu Q, Kurakake S, Okamura N, Kumae T, Umeda T, and Sugawara K.** Impact of a competitive marathon race on systemic cytokine and neutrophil responses. *Med Sci Sports Exerc* 35: 348-355, 2003.
118. **Suzuki K, Totsuka M, Nakaji S, Yamada M, Kudoh S, Liu Q, Sugawara K, Yamaya K, and Sato K.** Endurance exercise causes interaction among stress hormones, cytokines, neutrophil dynamics, and muscle damage. *J Appl Physiol* 87: 1360-1367, 1999.
119. **Suzuki M, Umeda T, Nakaji S, Shimoyama T, Mashiko T, and Sugawara K.** Effect of incorporating low intensity exercise into the recovery period after a rugby match. *Br J Sports Med* 38: 436-440, 2004.
120. **Trappe TA, Standley RA, Jemiolo B, Carroll CC, and Trappe SW.** Prostaglandin and myokine involvement in the cyclooxygenase-inhibiting drug enhancement of skeletal muscle adaptations to resistance exercise in older adults. *Am J Physiol Regul Integr Comp Physiol* 304: R198-205, 2013.
121. **Turner JE, Wadley AJ, Aldred S, Fisher JP, Bosch JA, and Campbell JP.** Intensive exercise does not preferentially mobilize skin-homing T cells and NK cells. *Med Sci Sports Exerc* 48: 1285-1293, 2016.
122. **Vella L, Markworth JF, Paulsen G, Raastad T, Peake JM, Snow RJ, Cameron-Smith D, and Russell AP.** Ibuprofen ingestion does not affect markers of post-exercise muscle inflammation. *Front Physiol* 7: 86, 2016.
123. **Vella L, Markworth JF, Peake JM, Snow RJ, Cameron-Smith D, and Russell AP.** Ibuprofen supplementation and its effects on NF-kappaB activation in skeletal muscle following resistance exercise. *Physiol Rep* 2: 2014.
124. **Verde T, Thomas S, and Shephard RJ.** Potential markers of heavy training in highly trained distance runners. *Br J Sports Med* 26: 167-175, 1992.
125. **Walsh NP, Gleeson M, Pyne DB, Nieman DC, Dhabhar FS, Shephard RJ, Oliver SJ, Berman S, and Kojanien A.** Position statement. Part two: maintaining immune health. *Exerc Immunol Rev* 17: 64-103, 2011.
126. **Walsh NP, Gleeson M, Shephard RJ, Gleeson M, Woods JA, Bishop NC, Fleshner M, Green C, Pedersen BK, Hoffman-Goetz L, Rogers CJ, Northoff H, Abbasi A, and Simon P.** Position statement. Part one: Immune function and exercise. *Exerc Immunol Rev* 17: 6-63, 2011.
127. **Wigernæs I, Hostmark AT, Stromme SB, Kierulf P, and Birkeland K.** Active recovery and post-exercise white blood cell count, free fatty acids, and hormones in endurance athletes. *Eur J Appl Physiol* 84: 358-366, 2001.
128. **Witard OC, Turner JE, Jackman SR, Kies AK, Jeukendrup AE, Bosch JA, and Tipton KD.** High dietary protein restores overreaching induced impairments in leukocyte trafficking and reduces the incidence of upper respiratory tract infection in elite cyclists. *Brain Behav Immun* 39: 211-219, 2014.
129. **Witard OC, Turner JE, Jackman SR, Tipton KD, Jeukendrup AE, Kies AK, and Bosch JA.** High-intensity training reduces CD8+ T-cell redistribution in response to exercise. *Med Sci Sports Exerc* 44: 1689-1697, 2012.



808

809 **Figure 1.** Overreaching or heavy training is associated with impaired cellular immune function

810 and latent viral reactivation. Reduction in lymphocyte trafficking and function during exercise

811 recovery impairs immune surveillance, which may increase susceptibility to opportunistic

812 infection. Lowered immunity may also allow previously acquired viruses to reactivate from a

813 latent state, which may cause further immunodepression and increase susceptibility to

814 infection.

815




Figure 2. Summary of the effects of nonsteroidal anti-inflammatory drugs (NSAIDs) and physical therapies on immune changes during recovery from exercise.

Table 1. Evidence heatmap comparing differences in immune responses to two versus one exercise bout on the same day (A), short vs long recovery between bouts on the same day (B), consecutive days of exercise (C), and weeks (D) or months (E) of intensified training.

	A			B			C			D			E		
Leukocyte cell count	5	3					2			1			2		
Neutrophil cell count	8	3		2			2	1		2			2		
Monocyte cell count	3	2			1					2			2	1	
Lymphocyte cell count	4	3	2		1		1			2			2		
Eosinophil cell count			1		1								1		
Oxidative burst activity			1		1	1	2								
Oxidative burst activity/neutrophil	1			1											
Plasma ELA concentration								1							
Plasma MPO concentration							1								
LPS-induced ELA release			1							1					
LPS-induced ELA release/neutrophil	1														
CD3 ⁺ T cell count	1		1				1						2		
CD4 ⁺ T cell count	4		2		1								2		
CD8 ⁺ T cell count	4	1	1	1									2		
CD4:CD8 ratio										1	1				
CD19 ⁺ B cell count	1	1											2		
CD4 ⁺ /CD69 ⁺ T cell count			1		1										
CD8 ⁺ /CD69 ⁺ T cell count			1	1											
Lymphocyte proliferation			2				1				1				
Antibody production		1									1				
Saliva IgA concentration	1	1								2				5	
Saliva IgA secretion rate	1	1					1	1						1	
CD16 ⁺ or CD56 ⁺ NK cell count	2	2		1			1			1			2		
CD56 ⁺ /CD69 ⁺ NK cell count	1				1										
NK cell cytotoxicity		1	1				1								
LPS-induced IL-6 production			1												
LPS-induced IL-8 production	1														
Leukocyte IL-8 mRNA								1							
Leukocyte IL-10 mRNA								1							
Leukocyte IL-1ra mRNA							1								
Plasma IL-6 concentration					1		2	1	1						
Plasma IL-1ra concentration					1		1								
Plasma IL-8 concentration							1								
Plasma IL-10 concentration							1								
Plasma TNF- α concentration							1		1						
Plasma MCP-1 concentration							1								
Muscle IL-6 mRNA expression							1								
Muscle IL-8 mRNA expression							1								
Muscle IL-1 β mRNA expression							1								
Muscle TNF- α mRNA expression							1								

Numbers represent the number of studies demonstrating an increase/greater change (red), no difference (green), or decrease/smaller change (blue) compared with the first bout of exercise, long recovery, before training, or healthy athletes (refer to text). Abbreviations: ELA, elastase; MPO, myeloperoxidase; NK, natural killer; MCP-1, monocyte chemoattractant protein 1

832

833